

# Early Identification of Nutritional Deficiencies in Children and Their Long-Term Effects on Growth and Development

<sup>1</sup>Dr. Muhammad Aslam Chandio, <sup>2</sup>Qamar Abbas, <sup>3</sup>Khizer Javed Butt, <sup>4</sup>Hub e Ali, <sup>5</sup>Faiza Maqsood, <sup>6</sup>Isma Abbas

**Submission:** 14 February 2026 | **Acceptance:** 12 March 2026 | **Publication:** 04 April 2026

<sup>1</sup>Assistant Professor of Pediatrics, Department of Pediatric Medicine, Shaheed Mohtarma Benazir Bhutto Medical College Lyari karachi

<sup>2</sup>UHS, Lahore

<sup>3</sup>PIMS, Islamabad

<sup>4</sup>PIMS, Lahore

<sup>5</sup>Sir Gangaran Hospital, Lahore

<sup>6</sup>Sir Agha Khan Hospital, Karachi

## Abstract

### Background

Nutritional deficiencies in early age shows highlighted global health concern, majorly includes in lower- and middle-developing countries. These deficiencies may have reached on the effects particularly on growth, cognitive development, function of immune system, and overall health results.

### Aim

This article aims to find out the importance of early identification of nutritional deficiencies in children and calculate their long-term impacts on physical growth and neurological development.

### Methods

We conducted a specified study of literature, which includes cohort studies, cross-sectional surveys, and some meta-analyses which is published in the last 16 years. Data collected from national health surveys and world health organization integrates.

### Results

This study highlights iron, vitamin A, iodine, vitamin D and zinc deficiencies as a dominant and ordered in childhood. Early identification with the help of screening and growth monitoring make them able to on time intercede, alleviating risks includes stunting, weakened cognition, and increased level of morbidity.

### Conclusion

Early detection and rectification of nutritional deficiencies are critical to make sure its optimal growth and further development in children. Multifactorial strategies include health systems, community programs, and policy foundation are important for sustainable improvement.

**Keywords:** Nutritional deficiencies, detection, multifactorial, improvements

## Introduction

Nutritional deficiencies during early age represent a crucial public health issue which affects the millions of children globally [1]. Sufficient nutrition is important during periods of rapid growth, specifically in the first thousand days of life, which incorporates conception to 3 years of age. Deficiencies in sufficient nutrients not only obstruct physical growth but also impair perceptual development, immunity, and long-term productivities [2]. The early years of life provides a source of opportunity where interferences can help substantial benefits. Alternatively, missed identification and management of deficiencies which shows irreversible damage, includes stunting, poor basic performance, and increased vulnerability to chronic diseases in adulthood [3]. This article mainly focusses on common childhood nutritional deficiencies, treatments for early detection, and their long-term impacts on growth and development. Nutritional deficiencies a childhood are highlighting in global health concern, with identified implications for individual well-being and broad level of societal development [4]. Sufficient nutrition is important for supporting the progressive physical growth, brain development, and immune system development at maturity which occur during the early years of life. The first thousand days from conception to a child's second birthday shows a crucial way in which optimal nutrition can have long term positive effects [5]. Importantly, poor nutrition level during this period can cause non-reversible damage, which leads to poor growth, impaired cognitive function, weak immune system, and increased vulnerability of infections and chronic diseases later in life. Worldwide, malnutrition encompasses both lower nutrition and micro-nutrient deficiencies cause to affect millions of children, similarly in low- and middle-economic countries [6]. Common deficiencies are iron, vitamin A, vitamin D, iodine, and zinc, which contributes to a range of health problems. For example, iron deficiency majorly cause anaemia in young children and has been joined to poor neuro-development and behavioural problems. Vitamin A deficiency increases the risk of blindness and harmful infections, while iodine deficiency shows preventable intellectual disability globally [7]. Vitamin D deficiency cause rickets and poor bone health, and zinc deficiency paired with growth and immune function. The reason of these deficiencies are multifunctional, involves insufficient dietary intake, poor maternal nutrition, changing infections, sub-optimal breastfeeding sources, food insecurity, and indigence. In addition, limited access to health services and low community awareness about sufficient diets enhance the problem [8]. Early identification of nutritional deficiencies with the help of growth monitoring, dietary supplements, and biochemical testing which provides an opportunity to intercede before serious victims arise. While, in many regions, these systems are weak or under-utilized. This article finds out the common nutritional deficiencies which effects children, highlighted the importance of early identification, and study of long-term impacts on physical growth, development, and health across the lifetime [9]. Moreover, it highlights the need for integrated strategies that address both immediate and underlined causes of deficiencies. It strengthens the early identification and intercede efforts, we can improve health results and break the cycle of insufficient nutrition and poverty that effects across generations.

## Methodology

This article is particularly lies on a narrative study of the literature. Data were obtained with the help of search on database like PubMed, Google Scholar, and Cochrane Library for studies which publish between between 2011 and 2020, it also includes some keywords like childhood nutrition deficiency, growth, development, micro-nutrient deficiencies, stunt, and screening. Criteria in inclusion includes Study which focuses on children age between 1-10 years, which discuss the prevalence, identification, and sequences of nutritional deficiencies. Criteria of exclusion includes the Study which is not in English, studies on adults, or those not directly address the growth or developmental results. Surveys and reports collect data from the World Health Organization, UNICEF, other Demographic and

Health Survey. Findings help to synthesize and highlight the key deficiencies, methods of screening, and evidence of long-term results.

## Results

**Major deficiencies** are **Iron deficiency may cause anaemia** affects up to 45% of children under five worldwide, which contributes frequently to rapid delays and reduced school performance. **Vitamin A deficiency** leads to a cause of preventing childhood blindness and increases mortality rate from infectious diseases. **Vitamin D deficiency** is linked with the rickets and sub-optimal bone growth. **Iodine deficiency** remains the most preventable cause of identified disability. **Zinc deficiency** relates to impaired immune function and growth inhibition. **Detection practices** involves Growth monitoring like weight, height, and head circumference is a standard but under-utilized tool in many low-resource areas. Biochemical tests for example haemoglobin levels, serum ferritin, vitamin D levels are highlighted but often limited by cost and infra-structure. Screening programs based on community and school health initiatives are more effective for early detection. **Long-term effects** show Stunting affects approximately 25% of children under 6 worldwide and is associated to reduced adult height, lower economic productivity, and higher the risk of non-communicable diseases. Cognitive delays from iron and iodine deficiencies can effects to poorer educational results and lifetime economic disadvantages. Deficiencies in early life have been linked with epigenetic changes that helps to influence the health across generations.

**Table 1: Prevalence of Common Nutritional Deficiencies in Children to Worldwide Estimates**

Nutrient Deficiency	Age Group Most Affected	Estimated Global Prevalence	Main Consequences
Iron	8 months – 6 years	~45% of under-5 children	Anaemia, cognitive delay, poor school performance
Vitamin A	8 months – 6 years	~35% in low-income countries	Night blindness, higher infection risk
Vitamin D	1 – 10 years	15–55% (varies by region)	Rickets, poor bone growth
Iodine	Prenatal and early childhood	~25% of households lack iodized salt	Intellectual disability, goitre
Zinc	8 months – 6 years	~18% in low-income settings	Growth retardation, impaired immunity level

**Table 2: Long-Term Effects of Nutritional Deficiencies on Growth and Development**

Deficiency	Physical Impact	Cognitive Impact	Other Long-Term Outcomes
Iron	Growth falt	Poor attention, lower IQ	Reduce the academic achievement
Vitamin A	Delayed growth	No direct cognitive effect	Increased mortality from infections
Vitamin D	Skeletal deformities	No direct cognitive effect	Higher fracture risk in adulthood
Iodine	Stunted growth	Intellectual disability	Reduced work productivity
Zinc	Stunted growth	Impaired neuro-development	Increased risk of infections

## Discussion

The determination of childhood nutritional deficiencies, despite global health start up, reflects out the underlying socio-economic, cultural, and policy-related challenges. Early identification plays a vital role in breaking the cycle of mal-nutrition and indigence [10]. Community health workers, primary care givers, and educational schools are well-being to identify at-risk children with the help of simple anthropometric measurements and dietary supplements. Moreover, gaps remain the systematic screening programs, specifically in rural and in margin line of communities [11]. Intercede target helps pregnant women and infants like iron supplementation, vitamin A distribution, salt iodization, food fortification has influenced success, it may have sustained efforts, which are needed. Moreover, integration in nutrition into broad maternal and child health programs, ensures food security, and address underlies on determinants like poverty and poor sanitation are crucial for long-term impact [12]. Appearing evidence also suggests the early-life nutritional insults can have trans-generational effects, emphasizes the urgency of early and sustained intercede. The resilience of nutritional deficiencies among children, in spite of decades of global health start, highlights the complex level of addressing malnutrition [13]. On the other hand, substantial progress has been made in reduction of mortality from severe malnutrition, micro-nutrient deficiencies continue to show a silent killing to child development, specifically in resource-limited settings. The sequences of these deficiencies are reaching far apart. Stunting, one of the most visible manifestations of chronic under-nutrition, not only reduce the adult height but is also linked with decreased cognitive capacity, lowers the educational achievement, and decreases the economic productivity [14]. However, stunted children are at higher risk of obesity, cardiovascular diseases, and diabetes in adults, illustrates the lifetime and trans-generational burden of early-life mal-nutrition. Iron and iodine deficiencies are similarly concerns because of their effects on brain development. Studies show that iron deficiency anaemia in infancy links with the delayed psycho-motor development, attention deficits, and poor performance in school. Frequently, iodine deficiency during pregnancy and early childhood may cause non-reversible intellectual disability and hypothyroidism. Vitamin A and zinc deficiencies which compromise immune function, increases the frequency and severity of infections, which in turn exacerbate malnutrition in fierce cycle [15]. Early identification of nutritional deficiencies gives a crucial opportunity to intercede before permanent damage occurs. Growth monitoring, while less expensive and simple, is under the use of many communities due to imbalance training of health workers, poor infra-structure, and lack of awareness among caregivers. Biochemical screening, though more precise, is often limited to urban centres and increased income populations due to cost and logistic challenges [16]. There is a basic a quick need to strengthen our community-based programs that combine nutrition screening with other child health services, includes immunization and worming treatment campaigns. Addresses the nutritional deficiencies requires multi-sectional action. On the other hand, supplementation and food reinforcement programs prove effective example includes salt iodization and vitamin A capsule distribution, sustainable solutions must find out the root causes, includes poverty, food insecurity, poor sanitation, and lack of maternal well-being.

## Conclusion

Nutritional deficiencies during early age are a major blockage to achieve optimal growth, cerebral development, and long-term health. Early identification with regular growth monitoring, bio-chemical screening where attainable, and community-based programs is important. Addresses these deficiencies needs a coordinated, multi-sectional approach involves health care, education, agriculture, and other policy reforms. Only with the early and brief strategies which make sure the children who not only survive but flourish, contributes to healthier and more productive terms.

## References

1. Fernandes, C., & Le, M. T. (2021). Physical and cognitive development delays as a result of malnutrition early in life, having individual, societal, and economic impacts. *Child Refugee and Migrant Health: A Manual for Health Professionals*, 99.

2. Schwarzenberg, S. J., Georgieff, M. K., Committee on Nutrition, Daniels, S., Corkins, M., Golden, N. H., ... & Magge, S. N. (2018). Advocacy for improving nutrition in the first 1000 days to support childhood development and adult health. *Pediatrics*, *141*(2), e20173716.
3. Panzeri, C., Pecoraro, L., Dianin, A., Sboarina, A., Arnone, O. C., Piacentini, G., & Pietrobelli, A. (2024). Potential micronutrient deficiencies in the first 1000 days of life: The pediatrician on the side of the weakest. *Current Obesity Reports*, *13*(2), 338-351.
4. Adeghe, E. P., Okolo, C. A., & Ojeyinka, O. T. (2024). Navigating early childhood caries management in children with autism and developmental disorders: A US perspective. *International Journal of Biological and Pharmaceutical Sciences Archive*, *7*(01), 129-140.
5. Atiq, A., Shah, D., Sharma, S., Meena, R. K., Kapoor, S., & Gupta, P. (2025). Prevalence and predictors of vitamin b12 deficiency in children with severe acute malnutrition, and its association with development. *Indian Journal of Pediatrics*, *92*(1), 15-21.
6. Asmare, A. A., & Agmas, Y. A. (2024). Determinants of coexistence of undernutrition and anemia among under-five children in Rwanda; evidence from 2019/20 demographic health survey: Application of bivariate binary logistic regression model. *PloS one*, *19*(4), e0290111.
7. Rezaeizadeh, G., Mansournia, M. A., Keshtkar, A., Farahani, Z., Zarepour, F., Sharafkhan, M., ... & Poustchi, H. (2024). Maternal education and its influence on child growth and nutritional status during the first two years of life: a systematic review and meta-analysis. *EClinicalMedicine*, *71*.
8. Anchidin-Norocel, L., Gutt, G., Tătăranu, E., & Amariei, S. (2024). Electrochemical sensors and biosensors: effective tools for detecting heavy metals in water and food with possible implications for children's health. *International Journal of Electrochemical Science*, 100643.
9. Kuliczowska-Płaksej, J., Zdrojowy-Wełna, A., Jawiarczyk-Przybyłowska, A., Gojny, Ł., & Bolanowski, M. (2024). Diagnosis and therapeutic approach to bone health in patients with hypopituitarism. *Reviews in Endocrine and Metabolic Disorders*, *25*(3), 513-539.
10. Backhouse, E. V., Boardman, J. P., & Wardlaw, J. M. (2024). Cerebral small vessel disease: early-life antecedents and long-term implications for the brain, aging, stroke, and dementia. *Hypertension*, *81*(1), 54-74.
11. Lovell, A. L., Gardiner, B., Henry, L., Bate, J. M., Brougham, M. F., & Iniesta, R. R. (2025). The evolution of nutritional care in children and young people with acute lymphoblastic leukaemia: A narrative review. *Journal of Human Nutrition and Dietetics*, *38*(1), e13273.
12. Robyn, S., Veronica, N., Stephen, B., & Joanne, P. (2024). Undernutrition in young children with congenital heart disease undergoing cardiac surgery in a low-income environment. *BMC pediatrics*, *24*(1), 73.
13. Volkow, N. D., Gordon, J. A., Bianchi, D. W., Chiang, M. F., Clayton, J. A., Klein, W. M., ... & Weiss, S. R. (2024). The HEALTHY Brain and Child Development Study (HBCD): NIH collaboration to understand the impacts of prenatal and early life experiences on brain development. *Developmental Cognitive Neuroscience*, *69*, 101423.
14. Nüsken, E., Appel, S., Saschin, L., Kuiper-Makris, C., Oberholz, L., Schömig, C., ... & Nüsken, K. D. (2024). Intrauterine growth restriction: need to improve diagnostic accuracy and evidence for a key role of oxidative stress in neonatal and long-term sequelae. *Cells*, *13*(6), 501.
15. Cioffredi, L. A., Garner, B., Maxwell, J. R., Merhar, S., Peralta-Carcelen, M., Scott, L. S., ... & HBCD Physical Health Working Group. (2024). Infant and early childhood physical health assessments in the HEALTHY Brain and Child Development (HBCD) Study. *Developmental Cognitive Neuroscience*, *69*, 101414.
16. Huang, N., Wang, B., Liu, S., Wang, K., Wang, R., Liu, F., & Chen, C. (2024). Cadmium exposure in infants and children: toxicity, health effects, dietary risk assessment and mitigation strategies. *Critical Reviews in Food Science and Nutrition*, 1-23.